

NSAM-913

CHANGES IN SPONTANEOUS ACTIVITY
AS A MEASURE OF SENSITIVITY TO
ROTATION IN THE WHITE RAT

Arnold Eskin and David C. Riccio

GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

Hard copy (HC) _____

Microfiche (MF) _____



JOINT REPORT

ff 653 July 65



FACILITY FORM 602

N65-35110

(ACCESSION NUMBER)

(THRU)

14

(PAGES)

1

(CODE)

CR 1-2253

(NASA CR OR TMX OR AD NUMBER)

04

(CATEGORY)

UNITED STATES NAVAL SCHOOL OF AVIATION MEDICINE
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

January 1965

CHANGES IN SPONTANEOUS ACTIVITY
AS A MEASURE OF SENSITIVITY TO
ROTATION IN THE WHITE RAT*

Arnold Eskin and David C. Riccio

Bureau of Medicine and Surgery
Project MR005.13-6001
Subtask 1 Report No. 103

NASA Order No. R-93

Approved by

Captain A. Graybiel, MC, USN
Director of Research

Released by

Captain H. C. Hunley, MC, USN
Commanding Officer

11 January 1965

*This research was conducted under the sponsorship of the Office of Life Science Programs, National Aeronautics and Space Administration.

U. S. NAVAL SCHOOL OF AVIATION MEDICINE
U. S. NAVAL AVIATION MEDICAL CENTER
PENSACOLA, FLORIDA

SUMMARY PAGE

THE PROBLEM

To investigate and measure the sensitivity of white rats to rotation using spontaneous activity as an index.

FINDINGS

Fifty-six unrestrained rats were individually exposed to a rotation speed between 0-18 RPM. Their activity was measured using a fourpoint scale: 0) no activity, 1) grooming and sniffing, 2) moderate running, and 3) rapid running. Amount of activity decreased as a function of rotation speed from 6 to 14 RPM, where it reached a lower limit plateau. Rate of decline within this speed range was also directly related to angular velocity. Postrotation activity was suppressed up to five minutes. The rats showed considerable sensitivity to Coriolis stimuli generated during rotation at constant angular velocities. A relationship was found between duration and magnitude of stimulation. These findings are encouraging for the use of behavioral methods in studying sensitivity to motion.

ACKNOWLEDGEMENTS

The authors wish to express their appreciation to Dr. Dietrich Beischer for his encouragement and support. The technical assistance of Mr. Calvin Crawford and Mrs. Beth Boss is gratefully acknowledged.

INTRODUCTION

The development of behavioral and physiologic measures of the effects of rotation on subhuman species for comparative purposes is a continuing problem in vestibular research. Predominantly physiologic symptoms such as nystagmic eye movements, sweating, pallor, vomiting, et cetera have been used widely to measure responses by man and animals to rotation (6,8,12,14,17). Although graded behavioral changes served to describe stages of canal sickness in squirrel monkeys and chimpanzees, vomiting was relied on to determine susceptibility thresholds to rotation (12,14).

Subhuman species like the white rat, which do not openly manifest these physiologic symptoms, have been considered as either insensitive to vestibular stimulation or refractory to practical measures of rotational effects. However, it is possible that these animals would show sensitivity to rotation through behavioral traits and that appropriate behavioral changes would function as a practical measure of sensitivity.

Since freely moving rats engage in a variety of spontaneous responses to their environment, e.g., from grooming and sniffing to rapid locomotion, it appeared that the animal's activity could provide the dependent variable for assessing rotation effects. The same variable has been utilized by Bindra (1) to assess the results of drugs on white rats. It was found that the physiological changes induced by the drugs could be sensitively measured by modifications in behavior patterns.

The purpose of this experiment was to investigate and measure the sensitivity of white rats to rotation using spontaneous activity as a behavioral index.

APPARATUS AND PROCEDURE

The subjects were 56 male albino Sprague-Dawley rats, three to four months of age. The animals were purchased in small lots from the Sprague-Dawley colony in Wisconsin and maintained in the Naval School of Aviation Medicine Laboratories.

The apparatus consisted of what might be termed a closed field for measuring activity. The closed field is intended to elicit a reasonable amount of spontaneous or exploratory activity from the rats, in contrast with the open field which characteristically reduces activity and produces other types of emotional behavior. The field was an enclosed area 2 feet square with 6-inch sides made of clear Plexiglas. A removable opaque cover with air holes reduced the illumination in the field and prevented the rat's escape. The closed field was mounted on a rotation turntable with speeds ranging from 0.17 to 28.0 RPM with a precision of ± 2 per cent. The height of the turntable was fixed to allow a seated observer convenient observation through the side of the field.

In order to investigate activity changes, five to eight randomly selected animals were individually tested at one of the following constant angular velocities: 0, 2, 4, ... 18 RPM. Each rat was used only once to avoid possible adaptation. The animals were run between 7:00 and 10:00 P.M. to take advantage of their increased night-time activity. The test room was dark except for a 50-watt red safety light located 5 feet from the rotation platform. The activity of each rat was measured during the prerotation, rotation, and postrotation periods. The duration of these three periods was six, six, and ten minutes, respectively. The postrotation session was longer to obtain data on the duration of the rotation effect. Three to four animals were run each evening, and at least two rotation conditions were employed to avoid any systematic bias. Three rats which froze continuously through the baseline period were discarded.

The activity of the rat was measured by observations taken systematically over time (cf. 2). The subject was observed and an activity score assigned for alternate ten-second periods. Thus, three activity scores, representing thirty seconds of observed behavior, were obtained each minute. The median activity per minute was then obtained for each group of rats at each rotation speed.

The scoring system was as follows:

- 3 - The highest level of activity. The rat ran rapidly and continuously in the field.
- 2 - The animal walked around, occasionally stopped and examined the field.
- 1 - The rat groomed, or scratched but remained fixed in one spot.
- 0 - Absolutely no movement observable. The rat held a "freeze" position.

The experimenter made an integrated judgment to assign a single score for a ten-second period. The use of half units (0.5, 1.5, et cetera) to modify the basic score provided a more finely graded continuum. Three personnel from the laboratory scored the rats at different times.

Thirty seconds after the rat was placed in the activity field, the first ten-second observation period began. After six minutes of baseline activity, the table was accelerated in two to three seconds to the predetermined rotation speed. Upon completion of this phase, the table was rapidly decelerated (two to three seconds) to 0 RPM. There was a thirty-second nonscored interval after starting and stopping rotation to permit angular acceleration effects to subside.

RESULTS

The six-minute prerotation period provided a baseline for each group. Since there were small differences in baselines of the different groups, the effect of various speeds on the animals is expressed directly as the ratio of activity during and after rotation to activity before rotation. Figure 1 presents the ratio of total rotation activity/total prerotation activity for all groups of subjects as a function of RPM. A smooth curve has been drawn through the experimental points. A slight increase in activity occurred at very low rotation speeds. Activity scores declined sharply after 6 RPM until reaching a plateau at about 14 RPM. The plateau shown in the figure after 14 RPM is the result of the scoring technique having reached its limit. In this range, the measure can no longer detect degrees of sensitivity, if there are any.

The time-course of activity changes for representative groups of animals is presented in Figure 2. Each point on the graph represents a group's median activity per minute. By the second minute of rotation exposure, animals at 10 RPM or above showed a sharp drop in activity rates. At speeds above 6 RPM the rate of activity decline was directly related to the rotation velocity. It can be seen in the figure that the control group showed a small decrease in activity during the twenty-two-minute session. The decline was most likely due to habituation to the box with an attendant reduction in exploratory responses.

It is clear from the curves in Figure 2 that both duration and speed of rotation were variables contributing to the animal's activity decline. This suggests the possibility of determining the various combinations of speed and exposure time which are equally severe as measured by activity change. Campbell and Kirby (3) have previously demonstrated that equal aversion contours can be plotted for a noxious stimulus such as electric shock. Figure 3 presents an "equal effect of rotation contour"; i.e., all points on the curve represent an equal reduction in activity. The points represent the time at which the activity of the different groups decreased by 30 per cent (reductions to other levels produce similar functions). The figure demonstrates, for example, that 1.3 minutes of rotation at 12 RPM produced a decrease in activity equivalent to three minutes of rotation at 8 RPM.

The animals rotated at 8 RPM or higher showed continued suppression of activity as long as five minutes after rotation. This can be seen in Figure 4 which presents the ratio of activities between the first five postrotation minutes and an equal portion of the baseline period as a function of the experimental conditions.

DISCUSSION

Unrestrained rats, like squirrel monkeys, show considerable sensitivity to constant angular velocities. Although vomiting was never observed in these rats, a characteristic "freezing" posture was noticed at higher velocities. The statue-like rigidity of the pose, readily discriminable from other behavior, has also been observed and referred to as "rotational posture" by Griffith (9) in his research on the effects of




Figure 1
Relative Activity as a Function of Angular Velocity




Figure 2

Time-Course of Activity for Representative Groups Prior To, During, and After Rotation




Figure 3

Duration of Rotation Exposure Required to Reduce Activity
by 30% at Various Rotation Speeds




Figure 4

Five Minute Postrotation Activity as a Function of Rotation Speed

rotation on nystagmus in the rat. That a laboratory animal as well studied and genetically controlled as the albino rat shows sensitivity to rotation suggests that it may be a useful subject for comparative vestibular research.

The question arises as to what factors are responsible for the reduction of activity. It is apparent from the slow time-course that the changes were not simply the result of the acceleration to the preset speed. A more likely variable was the Coriolis force which is a major source of vestibular input when the position of the head is moved relative to the rotating coordinate system of constant angular velocity (11). The Coriolis input is directly related to the velocity and magnitude of the subject's movement with respect to the rotating environment. Evidence presented in another study (15) demonstrated that the slight amount of G-forces present (maximum resultant G of 1.01) could not account for the observed changes and that the effects described were most likely attributable to Coriolis forces. The "Coriolis phenomenon" in man, which is of considerable importance for aspects of space medicine, has been the subject of intensive and continuing exploration by Graybiel and colleagues (4-7, 10, 13). In terms of labyrinthine mechanisms, Coriolis effects are presumably mediated by the semicircular canals rather than the otolith organs, hence the term "canal sickness" (6). By way of species comparison, canal sickness due to Coriolis acceleration has been reported for chimpanzees and squirrel monkeys (14).

The reason for the peculiar change in the postrotation activity suppression following exposure at 16 and 18 RPM shown in Figure 4 is not immediately clear. One possibility is that the animal was "thrown" into activity by the strong deceleration forces when the wheel was abruptly halted from high speeds. The animal could then quickly discriminate the lack of noxious Coriolis forces and resume its activity. Animals decelerated from slower speeds would have to emit several movements before learning that the aversive stimuli had been removed. Another possibility is that, due to the pronounced suppression of activity in the higher RPM groups, they were essentially recovering from the rotation effects before rotation ceased. The activity of the rotated animals recovered to the level of the control group after six to eight minutes.

The "equal effect of rotation curve" demonstrates that the behavioral response to rotation can be assessed as a joint function of duration and magnitude of rotation. The question arises as to why it took longer to obtain an equivalent activity decrease at 8 RPM than at 14 RPM. Coriolis input is generated by the rat when it moves in the rotating field. Accordingly, the postural freezing and reduction in activity can be considered as a learned response resulting from the reinforcement due to reduction of the Coriolis force which occurs when the head is held fixed. On the other hand, the decreased activity can also be viewed simply as a direct physiological response to "sickness" from rotation. Thus, the physiological effects of vestibular stimulation may accumulate within a certain period until they are equivalent to those produced by a 14 RPM stimulus over a briefer period. Alternatively, since the 8 RPM stimulus is less noxious than the 14 RPM, it may be that the more slowly rotated group was less motivated to learn to avoid the stimulation. This experiment does not discriminate among these possibilities and answer the question raised. The "equal effect contour" also

points out the fact that, after only three minutes of rotation at various speeds, the groups can be discriminated in terms of activity changes.

The small but consistent increase in activity at low angular velocities is difficult to explain. One possible interpretation is that it represents a transitory state which would eventually give way to cumulative Coriolis effects. To investigate this possibility, a group of rats was rotated at 4 RPM for fifteen minutes. The activity of the rotated group was found to remain slightly higher than a nonrotated control group for the same length of time. Within the explored time range and low rotation speed, the effects of rotation apparently were not cumulative.

To assess the generality of the activation phenomenon seen in the 4 RPM group, the relative activity data for the first minute of rotation was then examined for all nine groups (2 to 18 RPM). An initial increase in activity occurred in all but the high speed (12 to 18 RPM) groups for this period of time. At high rotation speeds, the Coriolis effects apparently occurred quickly enough to mask the initial response to rotation. At more moderate speeds (8 to 10 RPM), the amount of activity initially increased, then diminished during continued rotation exposure. Therefore, the increased activity of the rats seems to be a real phenomenon and may represent their attempt to escape an unusual vestibular stimulation.

These findings would seem to indicate that there is a certain Coriolis sensitivity threshold involved. It is difficult to assign an absolute threshold for this Coriolis sensitivity. At 7 RPM a 10 percent decrease occurred in baseline activity and at 9 RPM a 35 percent decrease for a five-minute rotation exposure. Therefore this range (7 to 9 RPM) represents a major transition in behavior. It is of interest that Riccio and Thach (16) using operant conditioning techniques have found that a major decrease in frequency of bar-pressing occurred between 10 and 14 RPM. In spite of a number of differences between the two experiments, both produced somewhat similar descriptions of the sensitivity of the rat to rotational forces.

The activity test used in this experiment has been found to provide a sensitive and quantitative index of rotational effects on rats. The system should therefore be useful for vestibular research. In addition, a long training period for the animals is not necessary. These features also recommend the system for its application in research concerned with pharmacological modification of sensitivity to motion sickness. The system has already proven useful in such an application by measuring the modification of rotation sensitivity through unilateral destruction of the hair cells of the labyrinth in rats by streptomycin sulfate injection (15).

REFERENCES

1. Bindra, D., and Baran, D., Effects of methylphenidylacetate and chlorpromazine on certain components of general activity. Jour. exp. anal. behav., 2: 343-350, 1959.
2. Bindra, D., and Blond, J., A time-sample method for measuring general activity and its components. Canad. J. Psychol., 12:74-76, 1958.
3. Campbell, B., and Kirby, R., Equal aversion contours for different types of electric shock stimuli. Paper presented at the 32nd Annual Meeting of the Eastern Psychological Association, April 7-8, 1961, Philadelphia, Pa.
4. Clark, B., and Graybiel, A., Human performance during adaptation to stress in the Pensacola Slow Rotation Room. Aerospace Med., 32:93-106, 1961.
5. Clark, R. T., Clamann, H. G., Balke, B., Tang, P. C., Fulton, J. D., Graybiel, A., and Vogel, J., Basic research problems in space medicine: A review. Aerospace Med., 31:553-577, 1960.
6. Graybiel, A., Clark, B., and Zarriello, J. J., Observations on human subjects living in a "slow rotation room" for periods of two days. Arch. Neurol., 3: 55-73, 1960.
7. Graybiel, A., Guedry, F. E., Johnson, W., and Kennedy, R., Adaptation to bizarre stimulation of the semicircular canals as indicated by the oculogyral illusion. Aerospace Med., 32:321-327, 1961.
8. Graybiel, A., Meek, J. C., Beischer, D. C., and Riopelle, A. J., Observations of canal sickness and adaptation in chimpanzees in a "slow rotation room." NSAM-456. Pensacola, Fla.: Naval School of Aviation Medicine, 1960.
9. Griffith, C. R., The effect upon the white rat of continued bodily rotation. Amer. Naturalist, 54:524-534, 1920.
10. Guedry, F. E., Kennedy, R., Harris, C. S., and Graybiel, A., Human performance during two weeks in a room rotating at three RPM. Aerospace Med., 35:1071-1082, 1964.
11. Guedry, F. E., and Montague, E. K., Quantitative evaluation of the vestibular Coriolis reaction. Aerospace Med., 32:487-500, 1961.
12. Johnson, W. H., Meek, J. C., and Graybiel, A., Effects of labyrinthectomy on canal sickness in squirrel monkeys. Ann. Otol., 71:289-298, 1962.

13. Kennedy, R., and Graybiel, A., Symptomatology during prolonged exposure in a constantly rotating environment at a velocity of one revolution per minute. Aerospace Med., 33:817-825, 1962.
14. Meek, J. C., Graybiel, A., Beischer, D. E., and Riopelle, A. J., Observations of canal sickness and adaptation in chimpanzees and squirrel monkeys in a "slow rotation room." Aerospace Med., 33:571-578, 1962.
15. Riccio, D. C., and Eskin, A., Some factors modifying the sensitivity of rats to rotation. In preparation.
16. Riccio, D. C., and Thach, J., Operant conditioning performance as a measure of the effects of rotation on rats. In preparation.
17. Wendt, G. R., Vestibular functions. In: Stevens, S. S. (Ed.), Handbook of Experimental Psychology. New York: John Wiley & Sons, Inc., 1951. Pp 1191-1223.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) U. S. Naval School of Aviation Medicine Pensacola, Florida		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE Changes in Spontaneous Activity as a Measure of Sensitivity to Rotation in the White Rat			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (Last name, first name, initial) Eskin, Arnold and Riccio, David C.			
6. REPORT DATE 11 January 1965		7a. TOTAL NO. OF PAGES 13	7b. NO. OF REFS 17
8a. CONTRACT OR GRANT NO. NASA Order No. R-93		9a. ORIGINATOR'S REPORT NUMBER(S) NSAM 913	
b. PROJECT NO. MR005.13-6001 Subtask 1		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) 103	
c.			
d.			
10. AVAILABILITY/LIMITATION NOTICES Copies of this report may be obtained from the Commanding Officer, U. S. Naval School of Aviation Medicine, Pensacola, Florida 32512 or from DDC.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
13. ABSTRACT Fifty-six unrestrained rats were individually exposed to a rotation speed between 0-18 RPM. Their activity was measured using a fourpoint scale: 0) no activity, 1) grooming and sniffing, 2) moderate running, and 3) rapid running. Amount of activity decreased as a function of rotation speed from 6 to 14 RPM, where it reached a lower limit plateau. Rate of decline within this speed range was also directly related to velocity. Postrotation activity was suppressed up to five minutes. The rats showed considerable sensitivity to Coriolis stimuli generated during constant speed of rotation. A relationship was found between duration and magnitude of stimulation. These findings are encouraging for the use of behavioral methods in studying sensitivity to motion. <i>author</i>			

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Spontaneous Activity Rotation Sensitivity Rat Behavioral Measurement Vestibular Stimulation Coriolis Acceleration Comparative Psychology						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

UNCLASSIFIED

Security Classification